

MIT-CTP-2728

hep-th/9805033

(2+1)-Dimensional QED, Anomalous Surface-Term Contributions and Superconductivity

Mu-In Park¹

Center for Theoretical Physics, Massachusetts Institute of Technology,
Cambridge, Massachusetts 02139 U.S.A.

ABSTRACT

It is shown that (2+1)-dimensional QED reveals several unusual effects due to the surface-term contributions. It is also shown that this system provides a new pairing mechanism for the high- T_c superconductivity on the plane.

PACS Nos: 11.10.Lm, 11.30.-j, 11.30.Cp, 74.70.Hk
20 June 1998

¹Electronic address: mipark@physics.sogang.ac.kr *and* mipark@ctp03.mit.edu

It is well known that (2+1)-dimensional gauge theories develop unusual features, due to the possibility of inclusion of the Chern-Simons (CS) term in the action [1]. Especially, due to this CS term, charged matter fields undergo unconventional transformation called “rotational anomaly” [2] by the improved Poincaré generators, constructed from the symmetric (Belinfante) energy-momentum tensor [3]. Furthermore, it was recently shown that the anomalous transformation is uniquely identified as the anomalous spin of the matter fields, which can be fractional, due to the unique meaning of the improved generators because they are gauge invariant on the constraints surface and obey the classical Poincaré algebra although this is not the case for the canonical (Noether) ones [4]. This fact is consistent with the fact that the rotational group in (2+1)-dimensions is $SO(2)$ which allows only continuous spectrum for the angular momentum. On the other hand, without doubt, the anomalous spin is nothing but the result of Aharonov-Bohm (AB) effect [5], since the CS term (in the Coulomb gauge) makes the charged matter fields carry the point magnetic vortices together with their electric charges which is the configuration of the AB effect. [This interpretation is still valid with the Maxwell term because only the asymptotic behavior is involved and this is controlled by the CS term [6].] So it has long been believed that this unusual rotational behavior is just because one has included the CS term which introduces the AB effect to the theory.

However, remarkably in a series of paper which are not well-known, Hagen showed that even the conventional QED in (2+1)-dimensions, (QED₃), also produces the rotational anomaly for the charged matter fields [7]. But there have been no further studies about the origin of this unexpected result. In this Letter, I investigate the physical and mathematical origin of this anomaly and relate this anomaly to the Aharonov-Casher (AC) effect [8] and/or the AB effect which is not manifest in the Lagrangian. Furthermore, I show that there is new anomalies in the time translation generated by Hamiltonian for the charged matter fields. Due to the uniqueness of the improved Poincaré generators, it is shown that this system also develops the fractional eigenvalue for the angular-momentum operator. This system shows also the attraction (repulsion) for the parallel (anti-parallel) magnetic dipole moments contrast to the usual (3+1)-dimensions case. It is shown that this system provides a new pairing mechanism for the high- T_c superconductivity on the plane.

The model to be studied is QED₃ with the massive relativistic complex scalars

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + (D_\mu\phi)^*(D^\mu\phi) - m^2\phi^*\phi, \quad (1)$$

where $g_{\mu\nu} = \text{diag}(1, -1, -1)$, $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ and $D_\mu = \partial_\mu + iqA_\mu$. (q is the electric charge of the elementary quanta of ϕ .) The equations of motion which follow from (1) are

$$\partial_\mu F^{\mu\nu} = J^\nu, \quad (2)$$

$$D_\mu D^\mu\phi + m^2\phi = 0 \quad (3)$$

and complex conjugate of (3). (J^ν is the conserved current $J^\nu = iq[\phi^* D^\nu \phi - (D^\nu \phi)^* \phi]$.) For the later purpose, let me first consider the (formal) solution of A^μ in (2). In order to make the gauge field configurations be rotationally symmetric, I choose the Coulomb gauge

$$\nabla \cdot \mathbf{A} = 0. \quad (4)$$

Then, from the temporal component of (2), $\nabla \cdot \mathbf{E} = J^0$ ($\mathbf{E} = -\nabla A^0 - \partial_t \mathbf{A}$), one obtains the instantaneous Coulomb potential and it's multipole expansion around $\mathbf{x}$

$$\begin{aligned} A^0(\mathbf{x}, t) &= \int d^2 \mathbf{x}' G(\mathbf{x} - \mathbf{x}') J^0(\mathbf{x}', t) \\ &= -\frac{1}{2\pi} Q \ln|\mathbf{x}| + \sum_{n=2}^{\infty} A_{(n)}^0(\mathbf{x}, t), \end{aligned} \quad (5)$$

where $G(\mathbf{x}) = -(1/2\pi) \ln|\mathbf{x}|$ is the two spatial-dimensions Green's function for the Laplacian, $\nabla^2 G(\mathbf{x}) = -\delta^2(\mathbf{x})$ and Q is the total electric charge $Q = \int d^2 \mathbf{x} J^0(x)$. [There is a constant ambiguity in the definition of $G(\mathbf{x})$, but this does not change the main issue of this Letter except in one minor point. This will be discussed later.] The direct Taylor expansion of $G(\mathbf{x}, t)$ gives the multipole expansion of the second line and $\sum_{n=2}^{\infty} A_{(n)}^0$ represent the (electric) dipole and higher-order moments terms and they behave as $\mathcal{O}(|\mathbf{x}|^{-1})$ at large $|\mathbf{x}|$. From the spatial components of (2), $\epsilon_{ij} \partial_j B - \partial_t E^i = J^i$ ($B = \nabla \times \mathbf{A} = \epsilon_{ij} \partial_i A^j$) one obtains the retarded vector potential and it's multipole expansion

$$\begin{aligned} A^i(\mathbf{x}, t) &= \int d^2 \mathbf{x}' G(\mathbf{x} - \mathbf{x}') [J_T^i(\mathbf{x}', t')]_{ret} \\ &= \frac{1}{2\pi} \frac{\mathbf{x}}{|\mathbf{x}|^2} \cdot \int d^2 \mathbf{x}' \mathbf{x}' [J_T^i(\mathbf{x}', t')]_{ret} + \sum_{n=3}^{\infty} A_{(n)}^i(\mathbf{x}, t), \end{aligned} \quad (6)$$

where $\mathbf{J}_T$ is the transverse current $J_T^i \equiv J^i - \partial_i \partial_t A^0 = -\epsilon_{ij} \partial_j \nabla \times \int d^2 \mathbf{x}' (1/2\pi) \ln|\mathbf{x} - \mathbf{x}'| \mathbf{J}(x')$, $\nabla \cdot \mathbf{J}_T = 0$ and $[f(\mathbf{x}', t')]_{ret} = f(\mathbf{x}', t - |\mathbf{x} - \mathbf{x}'|)$. $\sum_{n=3}^{\infty} A_{(n)}^i$ represents the expansion of quadrupole and higher-order moments terms and behaves as $\mathcal{O}(|\mathbf{x}|^{-2})$ at large $|\mathbf{x}|$. [The monopole term was neglected as usual by considering a well localized $[\mathbf{J}_T]_{ret}$ [9]] Now using the usual mathematical identity [8] $\int d^2 \mathbf{x}' (x^{i'} [J_T^j(x')]_{ret} + x^{j'} [J_T^i(x')]_{ret}) = 0$, (6) becomes

$$A^i(\mathbf{x}, t) = -\frac{1}{2\pi} \frac{\epsilon_{ij} x^j}{|\mathbf{x}|^2} [\mu](t; \mathbf{x}) + \sum_{n=3}^{\infty} A_{(n)}^i(\mathbf{x}, t), \quad (7)$$

where $[\mu] = (1/2) \int d^2 \mathbf{x}' \mathbf{x}' \times [\mathbf{J}_T(x')]_{ret}$ is the magnetic moment of the system induced by the retarded current $[\mathbf{J}_T(x')]_{ret}$. Note that $[\mu]$ has the implicit $\mathbf{x}$ -dependence through the retardation in $[\mathbf{J}_T(x')]_{ret}$ as well as the t -dependence. [These solutions (5)-(7) are only formal ones because one does not know the exact solution for J^μ in the classical as well as quantum

electrodynamics.] With these formal solutions, one gets B , E^i as

$$\begin{aligned} B(x) &= \frac{1}{2\pi} \nabla \cdot (\nabla \ln |\mathbf{x}| [\mu]) + \sum_{n=3}^{\infty} \nabla \times \mathbf{A}_{(n)}, \\ E^i(x) &= \frac{1}{2\pi} \frac{x^i}{|\mathbf{x}|^2} Q + \frac{1}{2\pi} \frac{\epsilon_{ij} x^j}{|\mathbf{x}|^2} [\dot{\mu}] + \sum_{n=3}^{\infty} [-\partial_i A_{(n-1)}^0 - \partial_t A_{(n)}^i], \end{aligned} \quad (8)$$

where over-dot represents the time derivative and higher-order moments terms of B and E^i behave as $\mathcal{O}(|\mathbf{x}|^{-3})$ and $\mathcal{O}(|\mathbf{x}|^{-2})$ at large $\mathbf{x}$, respectively.

Now, let me consider the canonical quantization with the usual commutation relations [2, 10]

$$\begin{aligned} [\phi(\mathbf{x}, t), \pi(\mathbf{x}', t)] &= [\phi^\dagger(\mathbf{x}, t), \pi^\dagger(\mathbf{x}', t)] = i\delta^2(\mathbf{x} - \mathbf{x}'), \\ [A^i(\mathbf{x}, t), E^j(\mathbf{x}', t)] &= -i\delta_{ij}^T(\mathbf{x} - \mathbf{x}'), \\ \text{others} &\text{ vanish,} \end{aligned} \quad (9)$$

with $\pi = (\partial_t - iqA_0)\phi^\dagger$, $\pi^\dagger = \phi(\partial_t + iqA_0)$, $\delta_{ij}^T(\mathbf{x}) = (\delta_{ij} - \partial_i \partial_j \nabla^{-2})\delta(\mathbf{x})$. [From now on, all quantum commutators are assumed to be the equal-time commutators.] To examine the Poincaré transformation and the covariance issues, let me consider the (manifestly) gauge invariant and symmetric energy-momentum tensor

$$T_s^{\mu\nu} = (D^\mu \phi)^*(D^\nu \phi) + (D^\nu \phi)^*(D^\mu \phi) + F^{\mu\rho} F_\rho^\nu - g^{\mu\nu} \left[(D_\mu \phi)^*(D^\mu \phi) - m^2 \phi^* \phi - \frac{1}{4} F_{\sigma\rho} F^{\sigma\rho} \right]. \quad (10)$$

From this, one can obtain the improved Poincaré generators, which are the gauge invariant constants of motion [3]

$$\begin{aligned} P_s^0 &= \int d^2\mathbf{x} \left[\pi\pi^\dagger + (\mathbf{D}\phi) \cdot (\mathbf{D}\phi)^\dagger + m^2 \phi\phi^\dagger + \frac{1}{2}(\mathbf{E}^2 + B^2) \right], \\ P_s^i &= \int d^2\mathbf{x} \left[\pi D^i \phi + (D^i \phi)^\dagger \pi^\dagger - \epsilon_{ij} E_j B \right], \\ M_s^{12} &= \int d^2\mathbf{x} \epsilon^{ij} x^i \left[\pi D^j \phi + (D^j \phi)^\dagger \pi^\dagger - \epsilon_{jk} E_k B \right], \\ M_s^{0i} &= tP_s^i - \int d^2\mathbf{x} x^i \left[\pi\pi^\dagger + (\mathbf{D}\phi) \cdot (\mathbf{D}\phi)^\dagger + m^2 \phi\phi^\dagger + \frac{1}{2}(\mathbf{E}^2 + B^2) \right], \end{aligned} \quad (11)$$

by $P_s^\mu = \int d^2\mathbf{x} T_s^{0\mu}$, $M_s^{\mu\nu} = \int d^2\mathbf{x} (x^\mu T_s^{0\nu} - x^\nu T_s^{0\mu})$. These are the well-known forms and produce the usual (first-order) Heisenberg equations and satisfy the Poincaré algebra in (3+1)-dimensions with the symmetric ordering. But as one shall see, this is not a trivial matter in (2+1)-dimensions due to the non-trivial surface term contributions. First of all, if one consider the spatial translation of the matter fields generated by P_s^i , it is easy to verify that one obtains the usual result $i[P_s^i, \phi(x)] = \partial^i \phi(x)$, where I have dropped the surface term

$\int d^2\mathbf{x}' \{ \partial'_i [(\nabla' \ln|\mathbf{x} - \mathbf{x}'|) \cdot \mathbf{A}(x')] - \nabla' \cdot [(\nabla' \ln|\mathbf{x} - \mathbf{x}'|) A^i(x')] - \nabla' \cdot [(\partial'_i \ln|\mathbf{x} - \mathbf{x}'|) \mathbf{A}(x')] \}$ because it vanishes as $|\mathbf{x}'|^{-1}$, $|\mathbf{x}'| \rightarrow \infty$ using the asymptotic behavior of $\mathbf{A}$ in (7). Next, if one consider the time translation generated by P_s^0 , one gets

$$i[P_s^0, \phi(x)] = \dot{\phi} - i \frac{q^2}{4\pi} \ln(\mathbf{0}) \phi - i \frac{q}{4\pi} \ln R|_{R \rightarrow \infty} [Q, \phi]_+, \quad (12)$$

where $[A, B]_+ = AB + BA$. The second term is a quantum correction arising from reordering at the same point [11] and is absent if one follows the old formulation of Pauli and Weisskopf without choosing the gauge [10]. Moreover this term appears only in the bosonic theory because of $(A^0)^2$ term in P^0 . The third term, which exists even in the classical level, is a surface contribution from $\int d^2\mathbf{x}' \nabla' \cdot (\mathbf{E}(x') \ln|\mathbf{x} - \mathbf{x}'|) = Q \ln R|_{R \rightarrow \infty}$ using the asymptotic behavior of $\mathbf{E}$ in (8). This is a kind of the infrared divergence which is independent on the mass of the matter field in (2+1)-dimensions. Here, it is noted that although (12) contains two unconventional terms, the second-order quantum equations of motion are the same form as the classical ones (3). This implies that the commutation relations (9) are a consistent choice. On the other hand, it is easy to see that the modified generator $\widetilde{P}_s^0 = \int d^2x \widetilde{T}_s^{00}$ with

$$\widetilde{T}_s^{00}(x) = T_s^{00}(x) - \left[\frac{q}{4\pi} \ln(\mathbf{0}) J^0(x) + \nabla \cdot \left(\frac{Q^2}{2(2\pi)^2} \frac{\mathbf{x}}{|\mathbf{x}|^2} \ln|\mathbf{x}| \right) \right] \quad (13)$$

gives the usual transformation $i[\widetilde{P}_s^0, \phi(x)] = \dot{\phi}$ but this is not allowed: The first term in the bracket (13) cancels the reordering effects in (12) and (14), but it breaks the Poincaré algebra involving $\widetilde{M}_s^{0i} = \int d^2\mathbf{x} (t T_s^{0i} - x^i \widetilde{T}_s^{00})$. Moreover, although the second term does not have this problem, $\widetilde{T}_s^{\mu\nu}$ does not transform as the second-rank tensor under the boost operator, i.e., $i[\widetilde{T}^{00}, M^{0i}] = (t \partial^i - x^i \partial_t) \widetilde{T}^{00} + 2T^{0i} - t(Q^2/8\pi^2) \partial^i \nabla \cdot (\frac{\mathbf{x}}{|\mathbf{x}|^2} \ln|\mathbf{x}|)$. These two facts are connected with the non-covariant forms of the vacuum expectation values of the the added terms in (13) [12]. Especially for the reordering term, it can be absorbed into the gauge transformation $\phi \rightarrow \phi e^{i\alpha t} \equiv \phi'$, $A_0 \rightarrow A_0 - \frac{\alpha}{q} \equiv A'_0$ with $\alpha = -(q^2/4\pi) \ln(\mathbf{0})$, but it is questionable that the fields ϕ' , which has no reordering effect, is meaningful one because it oscillates with infinite frequency. So, the two modification terms of (13) are not physical ones in these senses.

The Lorentz boost generated by M_s^{0i} becomes

$$\begin{aligned} i[M_s^{0i}, \phi(x)] &= (t \partial^i - x^i \partial_t) \phi(x) - i \frac{q}{2\pi} \int d^2\mathbf{x}' \ln|\mathbf{x} - \mathbf{x}'| [E^i(x') + (x^i - x^i) J^0(x'), \phi(x)]_+ \\ &+ \frac{i q^2}{4\pi} x^i \ln(\mathbf{0}) \phi(x). \end{aligned} \quad (14)$$

Here, I have neglected the surface contribution which comes from

$$\int d^2\mathbf{x}' \nabla' \cdot (x^i \mathbf{E}(x') \ln|\mathbf{x} - \mathbf{x}'|) = -Q \oint_{S_{R \rightarrow \infty}^1} d\theta \hat{\mathbf{r}}^i R \ln R \quad (15)$$

[The integration is evaluated on a circle with radius R , polar angle θ , and their corresponding orthogonal unit vectors $\hat{\mathbf{r}} = (\cos\theta, \sin\theta), \hat{\theta}$.] since it vanishes for any index i because of symmetry. However, note that the naive power counting gives $R \ln R|_{R \rightarrow \infty}$ divergence. So, in this case there is no surface contribution contrast to the naive expectation arising from the result (12). Returning to the result (14), the second term is the usual Lorentz transformation term for the matter field to accommodate with the Lorentz transformation of the gauge field $\mathbf{A}$ in the Coulomb gauge [13]. The third term is the reordering effect in the same origin with that of (12).

Finally, the rotation generated by M_s^{12} becomes

$$\begin{aligned} i[M_s^{12}, \phi(x)] &= \mathbf{x} \times \nabla \phi + i \frac{q}{2\pi} \int d^2 \mathbf{x}' \nabla' \times [\mathbf{A}(x') (\mathbf{x}' \cdot \nabla' \ln |\mathbf{x} - \mathbf{x}'|)] \phi(x) \\ &= \mathbf{x} \times \nabla \phi - i \frac{q}{2\pi} [\mu](t; |\mathbf{x}|)|_{|\mathbf{x}| \rightarrow \infty} \phi \end{aligned} \quad (16)$$

using the asymptotic form of $\mathbf{A}$ in (7) and the commutation relation $[[\mu], \phi] = 0$ which is implied by $[A^i, \phi] = 0$ in (9). Moreover, because of the fact that $[\mu](t; \mathbf{x})|_{|\mathbf{x}| \rightarrow \infty} = (1/2) \int d^2 \mathbf{x}' \mathbf{x}' \times \mathbf{J}_T(\mathbf{x}', t - |\mathbf{x}|)|_{|\mathbf{x}| \rightarrow \infty} \equiv \mu(t - \infty)$ and the anomalous term in (16) depends on the initial magnetic moment at the temporal infinity $t' = -\infty$, i.e., $\mu|_{t' = -\infty} \equiv \mu_{-\infty}$ for any finite time t . Here, there is no reordering effect in the same way as P_s^i but the surface contributions which were neglected in $[P_s^i, \phi]$ can not be neglected because of the existence of one more spatial coordinate in the integrand of M_s^{12} . This unconventional term which is fixed by the initial condition of the system at $t' = -\infty$, gives the anomalous spin for the matter field. This anomalous rotation transformation was previously noticed by Hagen [7] but the origin of this has been unclear. In order to understand the physical origin of the anomalous term more clearly, it is helpful to consider the surface contribution in the generator itself as

$$\begin{aligned} M_s^{12} &= M_c^{12} - \int d^2 \mathbf{x} \nabla \cdot [\mathbf{E} (\mathbf{x} \times \mathbf{A})] \\ &= M_c^{12} + \oint_{S_{R \rightarrow \infty}^1} d\theta R^2 \hat{\mathbf{r}} \cdot \mathbf{E} (\hat{\mathbf{r}} \times \mathbf{A}) \\ &= M_c^{12} + \frac{1}{2\pi} Q \mu_{-\infty} \end{aligned} \quad (17)$$

from which (16) is readily seen to follow: The first term is the canonical angular momentum operator $M_c^{12} = \int d^2 \mathbf{x} [\mathbf{x} \times (\pi \nabla \phi + (\nabla \phi)^* \pi^* - E_k \nabla A_k) + \mathbf{E} \times \mathbf{A}]$ which produces the usual angular momentum part and the second term gives the anomalous contribution of (16). Here, note that only $E_r = \hat{\mathbf{r}} \cdot \mathbf{E}$ and $A_\theta = \hat{\mathbf{r}} \times \mathbf{A}$ at the spatial infinity contribute to the surface term. But, from the asymptotic forms (7) and (8), only the field configurations $\mathbf{E}_{R \rightarrow \infty} = (1/2\pi)Q \hat{\mathbf{r}}/R$, $\mathbf{A}_{R \rightarrow \infty} = -(1/2\pi)\mu_{-\infty} \hat{\theta}/R$ are relevant [Since the next order terms in (7), (8) give the vanishing contribution $\mathcal{O}(R^{-1})$ to the surface term in (17) as $R \rightarrow \infty$.] and

they are the configurations for the plane projection of the infinite line charges (charge density Q per unit length) with a magnetic dipole moment $\mu = \mu_{-\infty}\hat{\mathbf{z}}$ in (3+1)-dimensions. This interpretation implies that the physical origin of the unconventional contribution in (16) is the AC effect [7, 14] (when the configuration is considered as that of the motion of magnetic dipole $\mu_{-\infty}$ in the electric field $\mathbf{E}_{R\rightarrow\infty}$) or the AB effect (when the configuration is considered as that of motion of charge Q in a magnetic field $\nabla \times \mathbf{A}_{R\rightarrow\infty}$). This fact is not manifest in the Lagrangian (1). Furthermore, since there is no reason for quantization of $\mu_{-\infty}$, the anomalous spin $s = -(1/2\pi)q\mu_{-\infty}$ can be fractional. This is the only known example of fractionally-valued total angular momentum in the parity-invariant gauge theory in (2+1)-dimensions without recourse to the CS term [15].

In order to give a unique meaning to the anomalous spin we must verify that the improved generators of (11) have more preferred meaning than the canonical ones. Usually, these two types of generators are identical on the constraint surface when one drops the surface term as in the (3+1)-dimensions case. But this is not true in (2+1)-dimensions in general; (17) is an example. Actually in this case the differences of these two types of generators, other than rotation operator, become

$$\begin{aligned} P_c^0 - P_s^0 &= \int d^2\mathbf{x} \nabla \cdot (\mathbf{E}A^0) = -\frac{1}{2\pi} \ln R|_{R\rightarrow\infty} Q^2, \\ P_c^i - P_s^i &= \int d^2\mathbf{x} \nabla \cdot (\mathbf{E}A^i) = 0, \\ M_c^{0i} - M_s^{0i} &= \int d^2\mathbf{x} \nabla \cdot [(tA^i - x^i A^0)\mathbf{E}] = 0, \end{aligned} \quad (18)$$

where I have used the asymptotic forms of (5)-(8) and the fact that geometrical term (15) vanishes. [If one allow the constant ambiguity in the Green's function as $G(\mathbf{x}) = -(1/2\pi)\ln|\mathbf{x}| + \text{const.}$, one can make $P_c^0 = P_s^0$ by choosing the constant as $(1/2\pi)\ln R|_{R\rightarrow\infty}$ but this has no effect in the physics, i.e., the anomalous contribution in (12). Actually this is the only modification produced by the constant ambiguity, which is only a minor point in our analysis] If one consider the (3+1)-dimensions, it is easy to verify that these surfaces terms all vanish as $P_c^0 - P_s^0 = \mathcal{O}(R^{-1})|_{R\rightarrow\infty}$, $P_c^i - P_s^i = \mathcal{O}(R^{-2})|_{R\rightarrow\infty}$, $M_c^{ij} - M_s^{ij} = \mathcal{O}(R^{-1})|_{R\rightarrow\infty}$ by power counting and $M_c^{0i} - M_s^{0i} = 0$ by the geometrical reason essentially the same as (15) [16]. So, the fact of inequivalence of the two types of the generators is a genuine effect in (2+1)-dimensions. A similar but not exactly the same things happen when there is the CS term instead of the Maxwell term [4]. Hence, in (2+1)-dimensions it seems that there are at least two inequivalent spin contents for the matter fields depending on the type of the angular momentum operator: One is integer from the canonical operator M_c^{12} and the other is fractional from the improved ones M_s^{12} . However, as can be verified by tedious calculations, it is not the canonical ones but the improved generators with the orderings in (11), which obey the

quantum Poincaré algebra: In the canonical case, the commutator of Lorentz boost becomes $[M_c^{0i}, M_c^{0j}] = -i\epsilon_{ij}(M_c^{12} + (1/2\pi)\mu_{-\infty}Q)$ and the anomalous term on the right-hand side can not be removed without modifying other generators. [This can be also easily verified by noting that the gauge invariant fields combinations $\mathbf{E}, B, \pi\pi^\dagger, (\mathbf{D}\phi) \cdot (\mathbf{D}\phi)^\dagger, \dots$ etc., only by which the Poincaré generators can be expressed, have no anomalous transformations and no gauge dependent Lorentz transformation term contrast to the charge bearing fields $\phi, \phi^\dagger$ and the gauge field $\mathbf{A}$. So the calculation with the gauge invariant fields should be the same as in (3+1)-dimensions case where the generators satisfy the usual Poincaré algebra.] From this fact the canonical generators can be discarded as the unphysical ones. Moreover, from the fact that the anomalous term in M_s^{12} can not be discarded by a redefinition of generators [2], the improved generators have a unique meaning consistently with Einstein's theory of gravity [17]. Hence the anomalous spin which comes only from M_s^{12} , has a unique meaning also. However, this uniqueness of anomalous spin does not imply the anomalous statistics, which has only artificial meaning similar to the CS gauge theory [4].

Finally, I discuss one more unusual phenomenon of our model and it's implication to the high- T_c superconductivity. If one imagine the non-relativistic identical point-particles (these need not be restricted to the bosons) with the same magnetic moment μ [I have omitted the retardation symbol [] because there is no retardation effect in this non-relativistic case], it gives an attractive δ -potential in the dipole approximation

$$U(\mathbf{x}) \cong -\mu B \cong -\mu^2 \delta(\mathbf{x}). \quad (19)$$

The oppositely oriented (i.e., opposite sign) magnetic dipoles are repulsive [18]. This is sharply contrast to the phenomena in (3+1)-dimensions: In (3+1)-dimensions, the corresponding potential for two identical dipoles is $U(\mathbf{x}) \cong -\mu \cdot \mathbf{B} \cong \mu \cdot \mu / (4\pi|\mathbf{x}|^3) - (2\pi/3)\mu \cdot \mu \delta^3(\mathbf{x})$ and so a collapse of the dipoles is avoided from the existence of infinite barrier represented by the first term; But there is no infinite barrier in (2+1)-dimensions case (19) such that the collapse is inevitable. Moreover, it is known that the quantum mechanics allows a stable bound state with the binding energy $\Delta = \epsilon / (e^{m/\mu^2} - 1)$ for the two-body system with the potential (19) [19]. [The stable N -body ground state can be also constructed and is correspond to a Hartree-Fock-type approximation.] Here, ϵ is the energy cut-off. Hence, this system provides a new pairing mechanism for the planer systems, different from the Cooper-pairing mechanism [20]: In this new mechanism, the “ferromagnetic order” is favored as in the recent *magnon exchange pairing mechanism* for the high- T_c *CuO* superconductors [21]. Actually, it is easy to compute a rough estimate of the necessary magnetic moment $\mu \cong \ln 2 \mu_B$ ($\mu_B = e/2m_e$ is the Bohr-magneton) in order to give a critical temperature $T_c \cong \Delta/k_B \cong 200 \text{ K}$ (k_B is the Boltzmann constant) between normal (non-binding) and superconducting (binding) phases. [Here, m and

ϵ are chosen as the electron mass m_e and its rest energy, respectively] The time-reversal and parity symmetries of the system are consistent with a recent experimental result [22].

In summary, I have shown that QED₃ reveals several unusual phenomena due to the surface term contributions. Now, let me address several comments. First, about the AC effect which was identified from the surface term in M_s^{12} , this can be also expected by noting that the matter field ϕ in the Coulomb gauge carries the radial electric field $\mathcal{E}(\mathbf{x}') = -(q/2\pi)(\mathbf{x} - \mathbf{x}')/(\mathbf{x} - \mathbf{x}')^2$ because of $[\mathbf{E}(\mathbf{x}', t), \phi(\mathbf{x}, t)] = \mathcal{E}(\mathbf{x}')\phi(\mathbf{x}, t)$, which is one needed constituent for the AC effect. And as one more necessary constituent, the magnetic moment is provided by the conserved current at $t' = -\infty$. One notable thing is that in the fermion case, the intrinsic magnetic moment of the fermion does not contribute additionally to the AB/AC effect through the surface term in (16) or (17): This is a completely analogous situation to the AC effect in CS gauge theory [2]. Moreover, if one includes the CS term in the Lagrangian (1), the AB effect, which comes from the CS term, appears rather than the AB/AC effect which comes from the Maxwell term: In this case the lower-derivative CS term will dominate the higher-derivative Maxwell term asymptotically [9, 23]. As for the superconductivity, it would be interesting to consider the finite temperature field theory to get a fully field theoretical description of the phenomena. As a final comment, although I have considered only the Coulomb gauge, the anomalous surface contributions, i.e., the anomalous field transformation and the anomalous spin are gauge independent: This is because the surface terms are gauge invariant for the rapidly decreasing gauge transformation function Λ under the gauge transformation $A_\mu \rightarrow A_\mu + \partial_\mu \Lambda$, $\phi \rightarrow \exp[-iq\Lambda]\phi$ and the commutator relation $[Q, \phi(x)] = -q\phi(x)$ is gauge independent [4].

I would like to thank Prof. S. Deser, C. R. Hagen, R. Jackiw, J. D. Jackson, I. Kogan for valuable conversations and informations and Prof. Y.-J. Park for warm hospitality. I also acknowledge the financial support of Korea Research Foundation made in the program year 1997. The present work is supported also in part by funds provided by the U.S. Department of Energy (D.O.E) under cooperative research agreement No. DF-FC02-94ER40818.

References

- [1] S. Deser, R. Jackiw, and S. Templeton, Phys. Rev. Lett. **48**, 975 (1982); Ann. Phys. (N.Y.) **140**, 372 (1982).
- [2] C. R. Hagen, Ann. Phys. (N.Y.) **157**, 342 (1984); Phys. Rev. **D31**, 2135 (1985); A. M. Polyakov, Mod. Phys. Lett. **A3**, 325 (1988); G. W. Semenoff, Phys. Rev. Lett. **61**, 517 (1988).
- [3] F. J. Belinfante, Physica (Utrecht) **7**, 449 (1940).
- [4] M.-I. Park and Y.-J. Park, “New Gauge Invariant Formulation of the Chern-Simons Gauge theory”, MIT-CTP-2702, hep-th/9803208 (1998).
- [5] W. Ehrenberg and R. E. Siday, Proc. Phys. Soc. London, Sec. **B62**, 8 (1949); Y. Aharonov and D. Bohm, Phys. Rev. **115**, 485 (1959).
- [6] R. Jackiw, Ann. Phys. (N.Y.) **201**, 83 (1990).
- [7] C. R. Hagen, Phys. Rev. Lett. **58**, 1074 (1987); Phys. Rev. **D36**, 3773 (1987).
- [8] Y. Aharonov and A. Casher, Phys. Rev. Lett. **53**, 319 (1984).
- [9] J. D. Jackson, *Classical Electrodynamics* (John Wiley and Sons, Inc., New York, 1962).
- [10] W. Pauli and V. Weisskopf, Helv. Phys. Acta, **7**, 709 (1934); J. D. Bjorken and S. D. Drell, *Relativistic Quantum Fields* (McGraw-Hill, Inc., New York, 1965).
- [11] R. Jackiw and S. Y. Pi, Phys. Rev. **D42**, 3500 (1990); M.-I. Park and Y.-J. Park, *ibid.*, **D50**, 7584 (1994).
- [12] D. G. Boulware and S. Deser, J. Math. Phys. **8**, 1468 (1967).
- [13] B. Zumino, J. Math. Phys. **1**, 1 (1960).
- [14] C. R. Hagen, Phys. Rev. Lett. **64**, 2347 (1990).
- [15] C. R. Hagen, Phys. Rev. Lett. **68**, 3821 (1992); M. E. Carrington and G. Kunstatter, Phys. Rev. **51**, 1903 (1995).
- [16] Recently, this kind of things in (3+1)-dimensions have been questioned by Lavelle and McMullan. But our explicit analysis shows that this is not true due to the geometrical reason; see M. Lavelle and D. McMullan, Phys. Rep. **279**, 1 (1997).

- [17] Motivated by the modified gravity theory as suggested by Callan *et al.*, one can consider the new improved generators, $\bar{P}^\mu = \int d^2\mathbf{x} \bar{T}^{0\mu}$, $\bar{M}^{\mu\nu} = \int d^2\mathbf{x} (x^\mu \bar{T}^{0\nu} - x^\nu \bar{T}^{0\mu})$ with $\bar{T}^{\mu\nu} = T^{\mu\nu} + c(g^{\mu\nu}\square - \partial^\mu\partial^\nu)f(\phi^*\phi)$. Here, c is constant and $f(\phi^*\phi)$ is some function of $\phi^*\phi$. But it is easy to verify that $\bar{P}^\mu = P^\mu$, $\bar{M}^{\mu\nu} = M^{\mu\nu}$, with reasonable assumptions a) no angle dependence of $f(\phi^*\phi)$ and b) $f(\phi^*\phi) \sim |\mathbf{x}|^{-\delta}$ ($\delta > 0$) as $|\mathbf{x}| \rightarrow \infty$. So, there is no change in the conclusion of this Letter; see C. G. Callan, S. Colman, and R. Jackiw, Ann. Phys. (N.Y.) **59**, 42 (1982).
- [18] A similar behavior was studied by Stern but exact identification remains to be clarified; see J. Stern, Phys. Lett. **B 265**, 119 (1991).
- [19] C. Thorn, Phys. Rev. **D19**, 639 (1979); K. Huang, *Quarks, Leptons and Gauge Fields* (World Scientific, Singapore, 1982); R. Jackiw, in M. A. M. Bég Memorial volume (World Scientific, Singapore, 1991).
- [20] L. N. Cooper, Phys. Rev. **104**, 1189 (1956).
- [21] G. Chen and W. A. Goddard III, Science **239**, 899 (1988).
- [22] S. Spielman *et al.*, Phys. Rev. Lett. **65**, 123 (1990).
- [23] K. Shizuya and H. Tamura, Phys. Lett. **B 252**, 412 (1990).